

**ANDERS HAMSTEN PUBLICATIONS 2008****Peer-reviewed articles**

Sjögren P, Sierra-Johnson J, Gertow K, Rosell M, Vessby B, de Faire U, **Hamsten A**, Hellénus M-L, Fisher RM. Fatty acid desaturases in human adipose tissue: relationships between gene expression, desaturation indexes and insulin resistance. *Diabetologia* 2008;51:328-335.

Boquist S, Ruotolo G, Skoglund-Andersson C, Tang R, Björkegren J, Bond MG, de Faire U, Brismar K, **Hamsten A**. Correlation of serum IGF-I and IGFBP-1 and -3 to cardiovascular risk indicators and early carotid atherosclerosis in healthy middle-aged men. *Clinical Endocrinology* 2008;68:51-58.

Wågsäter D, Björk H, Zhu C, Björkegren J, Valen G, **Hamsten A**, Eriksson P. ADAMTS-4 and -8 are inflammatory regulated enzymes highly expressed in macrophage-rich areas of human atherosclerotic plaques. *Atherosclerosis* 2008;196:514-522.

Sjögren P, Nordin Fredriksson G, Rosell M, de Faire U, **Hamsten A**, Nilsson J, Hellenius M-L, Fisher RM. Autoantibodies against modified apolipoprotein B-100 in relation to low-density lipoprotein size and the metabolic syndrome in otherwise healthy men. *Metabolism* 2008;57:362-366.

Broadbent HM, Peden JF, Lorkowski S, Goel A, Ongen H, Green F, Clarke R, Collins R, Franzosi MG, Tognoni G, Seedorf U, Rust S, Eriksson P, **Hamsten A**, Farrall M, Watkins H, for the PROCARDIS consortium. Susceptibility to coronary artery disease and diabetes is encoded by distinct, tightly linked, SNPs in the ANRIL locus on chromosome 9. *Human Molecular Genetics* 2008;17:806-814.

Skogsberg J, Lundström J, Kovacs A, Nilsson R, Noori P, Maleki S, Köhler M, **Hamsten A**, Tegner J, Björkegren J. Transcriptional profiling uncovers a network of cholesterol-responsive atherosclerosis target genes. *PLoS Genetics* 2008;4:e1000036.

Westerbacka J, Cornér A, Kolak M, Makkonen J, Turpeinen U, **Hamsten A**, Fisher RM, Yki-Järvinen H. Insulin regulation of MCP-1 in human adipose tissue of obese and lean women. *American Journal of Physiology (Endocrinology and Metabolism)* 2008;294:E841-845.

Mälärstig A, Eriksson P, Rose L, Diehl KA, **Hamsten A**, Ridker PM, Zee RYL. Genetic variants of tumor necrosis factor superfamily, member 4 and risk of incident atherothrombosis, and venous thromboembolism. *Clinical Chemistry* 2008;54:833-840.

Krapivner S, Popov S, Chernogubova E, Hellénus M-L, Fisher R, **Hamsten A**, van't Hooft FM. Insulin-induced gene 2 involvement in human adipocyte metabolism and body weight regulation. *Journal of Clinical Endocrinology and Metabolism* 2008;93:1995-2001.

Malarstig A, Sigurdsson S, Eriksson P, Samnegård A, Berne G, Hedin U, Wallentin L, Siegbahn A, **Hamsten A**, Syvänen A-C. Variants of the interferon regulatory factor 5 gene regulate expression of IRF5 mRNA in atherosclerotic tissue but do not confer susceptibility to myocardial infarction. *Arteriosclerosis, Thrombosis, and Vascular Biology* 2008;28:975-982.

Hägg DA, Jernås M, Wiklund O, Thelle DS, Fagerberg B, Eriksson P, **Hamsten A**, Olsson B, Carlsson B, Carlsson LMS, Svensson P-A. Expression profiling of macrophages from subjects with atherosclerosis to identify novel susceptibility genes. *International Journal of Molecular Medicine* 2008;21:697-704.

Greco D, Kotronen A, Westerbacka J, Puig O, Arkkila P, Kiviluoto T, Laitinen S, Kolak M, Fisher RM, **Hamsten A**, Auvinen P, Yki-Järvinen H. Gene expression in human NAFLD. *American Journal of Physiology (Gastrointestinal and Liver Physiology)* 2008;294:G1281-1287.

Sevastianova K, Sutinen J, Kannisto K, **Hamsten A**, Ristola M, Yki-Järvinen H. Adipose inflammation and liver fat in patients with highly active antiretroviral therapy-associated lipodystrophy. *American Journal of Physiology (Endocrinology and Metabolism)* 2008;295:E85-91.

Mälarstig A, Eriksson P, **Hamsten A**, Lindahl B, Wallentin L, Siegbahn A. Raised interleukin-10 is an indicator of poor outcome and enhanced systemic inflammation in patients with acute coronary syndrome. *Heart* 2008;94:724-729.

Sjögren P, Fredriksson GN, Samnegård A, Ericsson C-G, Fisher RM, Nilsson J, **Hamsten A**. High plasma concentrations of autoantibodies against native peptide 210 of apoB-100 are related to less coronary atherosclerosis and lower risk of myocardial infarction. *European Heart Journal* 2008;29:2218-2226.

Skogsberg J, Dicker A, Rydén M, Åström G, Nilsson R, Bhuiyan H, Vitols S, Mairal A, Langin D, Alberts P, Walum E, Tegner J, **Hamsten A**, Arner P, Björkegren J. The LDL receptors on adipocytes mediate inhibition of lipolysis by binding to apolipoprotein B-containing LDL. *PLoS One* 2008;3:e3771.

**Hamsten A**, Eriksson P. Identifying the susceptibility genes for coronary artery disease: from hyperbole through doubt to cautious optimism. *Journal of Internal Medicine* 2008;263:538-552.