

MARIE WAHREN-HERLENIUS PUBLICATIONS 2008**Peer-reviewed articles**

Hennig J, Bresell A, Sandberg M, Hennig KD, **Wahren-Herlenius M**, Persson B, Sunnerhagen M. The fellowship of the RING: the RING-B-box linker region interacts with the RING in TRIM21/Ro52, contains a native autoantigenic epitope in Sjögren syndrome, and is an integral and conserved region in TRIM proteins. *J Mol Biol*, 2008 377:431-49.

Skog A, **Wahren-Herlenius M**, Sundström B, Bremme K, Sonesson SE. Outcome and growth of infants fetally exposed to heart block-associated maternal anti-Ro52/SSA autoantibodies. *Pediatrics*, 2008 121:e803-809.

Strandberg L, Ambrosi A, Espinosa A, Ottosson L, Eloranta ML, Zhou W, Elfving A, Greenfield E, Kuchroo VK, **Wahren-Herlenius M**. Interferon- α induces up-regulation and nuclear translocation of the Ro52 autoantigen as detected by a panel of novel Ro52-specific monoclonal antibodies. *J Clin Immunol*, 2008, 28:220-31.

Popovic K, **Wahren-Herlenius M**, Nyberg F. Clinical follow-up of 102 anti-Ro/SSA-positive patients with dermatological manifestations. *Acta Derm Venerol* 2008 88(4):370-5.

Strandberg L, Winqvist O, Salomonsson S, Mohseni S, Buyon J, Julkunen H and **Wahren-Herlenius M**. Antibodies to amino acid 200-239 (p200) of Ro52 as serological markers for the risk of developing congenital heart block. *Clin Exp Immunol* 2008, 154:30-7.

Espinosa A, Oke V, Elfving A, Nyberg F, Covacu R., **Wahren-Herlenius M**. The autoantigen Ro52 is an E3 ligase resident in the cytoplasm but enters the nucleus upon cellular exposure to nitric oxide. *Exp Cell Res* 2008, 314:3605-13.